



Área: Fonoaudiologia 1207

PREVALENCE OF ENURESIS IN CHILDREN WITH CLEFT LIP AND PALATE: PRELIMINARY RESULTS

Fernandes MBL*3, Trindade IE

From the Laboratory of Physiology, Hospital for Rehabilitation of Craniofacial Anomalies (HRAC), University of São Paulo, Bauru, Brazil.

OBJECTIVES: Nocturnal enuresis (NE) is a common childhood condition, present in about 15% of Brazilian children. There is ongoing debate regarding the relationship between obstructive sleep apnea syndrome (OSAS) and NE. Craniofacial anomalies predispose to upper airway obstruction and OSAS. To study: (1) the prevalence of nocturnal enuresis (NE), diurnal urinary incontinence (DI) and OSAS in children with unilateral cleft lip and palate (UCLP) nonsyndromic, (2) risk factors associated with NE. METHODS: Here are preliminary results of an observational cross-sectional study of 83 children (54% males) 6-12 years of age (mean age 10.4 ± 1.82 years) with UCLP nonsyndromic. Study consisted of a personal interview with the child/caregivers and chart review. Severity of NE and DI, gender, tonsillectomy and/or adenoidectomy, frequency, arousal and sleeping disturbances were assessed. OSAS was identified by the presence of snoring, intermittent pauses and/or gasps. The Sleep Disturbance Scale for Children (SDSC) cut point sleep-disordered breathing (SDB>6) for OSAS. RESULTS: Nocturnal enuresis was reported in 15 of 83 children (18%) (8 males, mean age 9.6±1.32 years), and of these 4(27%) had associated DI. Twelve (84%) had mild enuresis (1-2nights/week). Thirty-four children (40%) had SDB>6.Among 17(21%) children who had previously undergone tonsillectomy and/or adenoidectomy (T&A) 3 (17%) had NE. Of the 15(18%) children with NE 20% had undergone T&A in the past. Five out of 15 enuretic children (30%) had symptoms suggestive of obstructive sleep apnea syndrome (SDB>6). CONCLUSION: Almost one fourth of children UCLP nonsyndromic had enuresis. OSAS was present in about one third of enuretic children. Tonsillectomy and/or adenoidectomy did not affect the prevalence of enuresis.