

NASAL OBSTRUCTION INCREASES THE RISK OF OBSTRUCTIVE SLEEP APNEA?

Fernandes MBL*, Trindade IEK

From the Laboratory of Physiology, Hospital for Rehabilitation of Craniofacial Anomalies (HRAC), University of São Paulo, Bauru, Brazil.

OBJECTIVE: Craniofacial anomalies predispose to upper airway obstruction. Obstructive sleep apnea syndrome (OSAS) is related obesity, hypertension, attention deficit and learning, nocturnal enuresis. **TO STUDY:** prevalence and association of nasal obstruction and OSAS. **METHODS:** Here are preliminary results of an observational cross-sectional study of 83 children (54% males) 6-12 years of age (mean age 10.4 ± 1.82 years) with unilateral cleft lip and palate (UCLP) nonsyndromic. Study consisted of a personal interview with the child/caregivers. Congestion Quantifier Five-Item Test (CQ5) for nasal, patient with score of ≤ 6 are at a level that warrants examination and possible treatment. SN-5 survey as a measure of longitudinal change in health related quality of life (HRQoL). Visual Analog Scale (VAS), a child was asked to evaluate the level of the obstruction of his/her nose. OSAS was identified by the presence of snoring, intermittent pauses and/or gasps. The Sleep Disturbance Scale for Children (SDSC) cut point sleep-disordered breathing ($SDB > 6$) for OSAS. **RESULTS:** Twenty-nine children (35%) presented with CQ5 ≤ 6 . Mean SN-5 score was $1.8 (\pm 1.97)$. Mean SDB $6.3 (\pm 2.94)$. Thirty-four children (40%) had $SDB > 6$ (mean 9.3 ± 3.01). At baseline, the mean VAS on the cleft side was $5.8 (\pm 3.13)$ and noncleft side was $9.1 (\pm 3.52)$. Symptoms of obstructive sleep apnea syndrome (OSAS) with $SDB > 6$ were observed in 69% of children with CQ5 ≤ 6 (mean 11.8 ± 5.92). **CONCLUSION:** Children with nonsyndromic UCLP present high prevalence of symptoms suggestive of obstructive sleep apnea syndrome (OSAS). Symptomatic nasal obstruction increases incidence of symptoms of OSAS.