O IMPACTO DA SAÚDE BUCAL NA QUALIDADE DE VIDA DE PACIENTES COM FISSURAS LÁBIO-PALATINAS DO CAIF-CURITIBA

BALDERRAMA ÍF***[1], WOBETO APP[2], SILVA MAR[2], ALANIS LRA[2], SILVA JYB[3], WERNECK RI[2]

1. Mestranda em Periodontia na Faculdade de Odontologia de Bauru-Universidade de São Paulo, Bauru, Brasil
2. Pontifícia Universidade Católica do Paraná, Curitiba, Brasil
3. Centro de Atendimento Integral de Fissurado Lábio Palatal, Curitiba, Brasil

OBJETIVO: Avaliar o impacto da fissura lábio-palatina na qualidade de vida dos pacientes do Centro de Atendimento Integral ao Fissurado Lábio Palatal de Curitiba - PR (CAIF). **MÉTODOS:** Para mensurar a interferência da saúde bucal na qualidade de vida foram utilizados indicadores sócio-dentais que avaliam dimensões físicas, psicológicas e sociais e são baseados na autopercepção do indivíduo e nos impactos odontológicos sofridos. A coleta de dados foi realizada utilizando o questionário IODD (Índice de Impactos Odontológicos no Desempenho Diário) para determinar a qualidade de vida em pacientes com fissuras lábio-palatinas do CAIF. Os prontuários foram analisados em relação à identificação do paciente, classificação e lado de ocorrência da fissura. **RESULTADOS:** Os 231 pacientes entrevistados possuíam idade variando de 7 a 65 anos (média: 19,75). Dentre estes, 44,6% eram do sexo feminino e 55,4% do sexo masculino. Em relação à prevalência dos tipos de fissura, a fissura transforame incisivo unilateral esquerda foi a mais prevalente. Em relação ao questionário IODD, este foi utilizado com o intuito de compreender a quantidade de pacientes que responderam que a sua fissura o afeta em determinada atividade. Utilizando o teste qui-quadrado, foi observado que o índice com maior impacto na qualidade de vida foi "falar e pronunciar com clareza, quando associamos a severidade da fissura e a atividade (p= 0,008). Já com o teste t-student foi verificado relação entre qualidade de vida para "dormir e relaxar" e pacientes com menos cirurgias que a média (5,01 cirurgias) (p=0,031). **CONCLUSÃO:** Podemos afirmar que existe uma relação entre as fissuras lábio-palatinas e qualidade de vida.

Apoio Financeiro: Bolsa PUCPR-PIBIC

THE ORAL HEALTH IMPACT ON THE QUALITY OF LIFE IN PATIENTS WITH CLEFT LIP AND PALATE OF CAIF - CURITIBA

BALDERRAMA ÍF***[1], WOBETO APP[2], SILVA MAR[2], ALANIS LRA[2], SILVA JYB[3], WERNECK RI[2]

1. DDS, MS Student, Discipline of Periodontics, Department of Prosthodontics, Bauru School of Dentistry, University of São Paulo, Bauru, SP, Brazil
2. Pontifical University Catholic of Parana, Curitiba, Brazil
3. Cleft Lip and Palate Integral Care Center- CAIF, Curitiba, Brazil

BACKGROUND: *The aim of this study is to evaluate the influence of cleft lip and palate on the quality*

of life in patients at the Cleft Lip and Palate Integral Care Center (CAIF) - Curitiba, PR, Brazil. **METHODS:** *To measure the interference of oral health on the quality of life, are used socio-dental indicators that evaluate physical, psychological and social dimensions and are based on the self-awareness of the individual and in dental impacts suffered. Data collection was performed using the questionnaire IODD (Index of the Oral Impacts on Daily Performance) to measure quality of life of 115 patients with cleft lip and palate at the CAIF. Record of these patients was performed through to patient identification, classification and occurrence of the cleft.* **RESULTS:** *231 patients interviewed ages from 7-65 years (mean: 19,75). Of these, 44,6% were female and 55,4% male. The most frequent cleft type was unilateral left trans-incisive foramen cleft. The questionnaire IODD used with aim for understanding the amount patients who responded that your cleft affects in your particular activity. The chi-squared test statstic was observed the index with the greatest impact "speaking and pronounce clearly" when associate the severity of the cleft and activity (p=0,008). With student's-t test statistic was checked relationship between quality of life for "sleeping and relaxing" and patients with fewer surgeries that mean (5,01 surgery) (p=0.031).* **CONCLUSION:** *There exist a relationship between the cleft lip and palate and quality of life.*

Financial Support: Scholarship PUCPR-PIBIC